


INSTRU

Concentrated disinfectant for critical medical and dental instruments

HISEPT INSTRU is a concentrated disinfectant used for disinfection of critical medical and dental Instruments before sterilization. Highly effective antimicrobial formulation offers broad-spectrum activity even against mycobacterium and non-enveloped viruses. This product has a concentrated formulation and due to the presence of corrosion inhibitor compounds, is compatible with most materials. Aldehyde-free nature of HISEPT INSTRU prevents protein fixation.



Characteristics

- Broad-spectrum antimicrobial activity
- Good cleaning performance
- Cost-effective concentrated product
- Aldehyde and phenol free
- Usable in ultrasonic baths
- Reusable up to 14 days (depending on the level of pollution)

Product Specification

Color/Form	Colorless to pale yellow clear solution
pH	7-10
Scent	Characteristic odor
Shelf Life	24 months
Solubility (in water)	Completely soluble



THIS IS A HISEPT PRODUCT, WITH
**HIGH-LEVEL
SAFETY &
PROTECTION**

Ingredients

100 g HISEPT INSTRU contains:

- 3.8 g Polyhexamethylene biguanide hydrochloride
- 5.6 g Didecyldimethylammonium chloride

Auxiliary ingredients:

- Surfactants
- Corrosion inhibitors
- Solubilizer
- Dye
- Deionized water

Microbiological Effectiveness

Meets European Standards

Bactericidal Activity

According to EN 13727 Staphylococcus aureus/ ATCC 6538
(contact time: 15 minutes Escherichia coli/ ATCC 10536
use dilution: 2%) Pseudomonas aeruginosa / ATCC 15442
Enterococcus hirae/ ATCC 10541

According to EN 14561 Staphylococcus aureus/ ATCC 6538
(contact time: 15 minutes Pseudomonas aeruginosa / ATCC 15442
use dilution: 2%) Enterococcus hirae/ ATCC 10541

Yeasticidal Activity

According to EN 13624 Candida albicans/ ATCC 10231
(contact time: 15 minutes
use dilution: 2%)

According to EN 14562 Candida albicans/ ATCC 10231
(contact time: 15 minutes
use dilution: 2%)

Fungicidal Activity

According to EN 13624 Aspergillus niger/ ATCC 16404
(contact time: 15 minutes
use dilution: 2%)

According to EN 14652 Aspergillus niger/ ATCC 16404
(contact time: 30 minutes
use dilution: 2%)

Virucidal Activity

According to EN 14476 Herpes simplex virus(HSV-1)
(contact time: 15 minutes
use dilution: 2%)



Directions for use

The concentrate shall be diluted before use with water to 2%. Immerse clean devices completely into diluted solution for at least 15 minutes.

Compatibility

HISEPT INSTRU is compatible with most materials such as stainless steel, copper, plastic, rubber and silicon.

2/2 Latest revision: Dec 1, 2023

Packagings

3 canisters per carton,	5 L each
6 bottles per carton,	1 L each

Hazard Rating

Concentrate	NFPA
Health	1
Flammability	3
Reactivity	1



Warning

- The disinfection tank shall always be covered with a lid. If the solution is kept in a closed container, it will be reusable for up to 14 days.
- Hazardous in case of skin and eye contact, ingestion or inhalation.
- Keep out of reach of children and unaware persons.
- Store between + 5 °C and + 35 °C and away from sunlight.



Certifications

- >> ISO 9001: 2015
- >> ISO 13485: 2016
- >> GMP
- >> FSC



HISEPT[®]
HIGH-LEVEL SAFETY & PROTECTION

Middle East Office
Behban Pharmed Lotus Co.
Address : Unit 113, 12 Floor, Bldg. B, Kaveh
Complex, Azadi Ave., Tehran, Iran
Tel: (+9821) 66581294-5/66581269

info@hisept.com

World Locations: Düsseldorf, Kuala Lumpur, Istanbul, Tehran, Baghdad